



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

An Enumeration of the Plants Collected by Dr. H. H. Rusby in  
South America, 1885-1886.—X.

(Continued from Vol. XVI, p. 327.)

*Inga strigillosa*, Spruce in Mart. Fl. Bras. Reis, 1,500 ft. (1002).

*Inga nobilis*, Willd. Enum. 1047. Mapiri, 2,500 ft. (1003).

*Inga edulis*, Mart. Herb. fl. Bras. 113. Unduavi, 8,000 ft. (991); Junction of the Beni and Madre de Dios (995).

*Inga marginata*, Willd. Sp. Pl. iv. 1015. Guanai, 2,000 ft. (994; 998; 999).

*Inga punctata*, Willd. l. c. 1016. Beni River, (1000).

*Inga Matthewsiana*, Benth. in Hook. Lond. Journ. Bot. iv. 594. Guanai, 2,000 ft. (996).

*Inga tomentosa*, Benth. Trans. Linn. Soc. xxx. 609. Mapiri, 5,000 ft. (997).

*Inga stipularis*, DC. Mem. Leg. 440. Mapiri, 5,000 ft. (1004).

INGA BOLIVIANA, spec. nova. Flores tomentelli, in receptaculo distincte pedicellati, umbellam globosam formantes; pedicelli calyce longiores; petiolus nudus; foliola 3-4 juga, elliptico-oblonga, apice acuminata, utrinque glabra, venulosa, sessiles, 9-20 cm. longa; umbellæ longiuscule pedunculatæ, pedunculi tomentelli; pedicelli 5-8 mm., calyc. 4-6 mm., corolla 10 mm. longa; legumen ignotum. Junction of the Rivers Beni and Madre de Dios (990). This is nearest to *I. quaternata*, Poepp. & Endl. but has much larger leaflets and less woolly flowers.

*Inga*, probably undescribed; collected only in fruit. Junction of the Beni and Madre de Dios (993 and 2375). Some flowers of *I. edulis*, Mart. were accidentally distributed with No. 993.

*Inga*, related to *I. nobilis*, but with thin membranaceous leaves and simple inflorescence. Mapiri, 2,500 ft. (1001). This is most likely an undescribed species.

ROSACEÆ.

*Licania Benthami*, Hook. f. in Mart. Fl. Bras. xiv. (2) 12. Junction of the Madre de Dios and Beni (2420). The same as Spruce No. 3278.

LICANIA PALLIDA, spec. nova. Folia oblonga, utrinque pallida et glabra, 10-12 cm. longa, 5 cm. lata; petiola crassa, 1 cm. longa; paniculæ terminales et axillares; ramulis, bracteolis, pedunculis calicibusque albo-pubescentibus; flores spicati;

spicæ 3-4 cm. longæ, 8mm. latæ; pedunculi 2 mm. longi; bracteolæ longiores; stamina sub 5, exserta, petala longiores; fructus ignotus. Junction of the Beni and Madre de Dios (2442). Apparently nearest to *L. polita*.

*Hirtella Americana*, Aubl. Guian. i. 247. Guanai, 2,000 ft. (702); Mapiri, 5,000 ft. (704). Junction of the Beni and Madre de Dios (1371). A widely diffused and variable species.

*Hirtella bullata*, Benth. in Hook. Journ. Bot. ii. 216. Reis, 1,500 ft. (703). The same as Spruce No. 3084, Herb. Kew.

*HIRTELLA BURCHELLII*, spec. nova. Flores in paniculas angustas dispositi; paniculæ rufo-villosæ, angustæ, axillares terminalibusque, 12-20 cm. longæ; ramulis rufo-villosis; foliis ovato-oblongis, acuminatis, basi rotundatis, utrinque nitidis, supra lævis, subtus ad nervos parce villosis; pedunculis 2mm. longis calicibusque dense villosis. Beni River (1222). The same as Burchell's Nos. 6331, 6416 and 6571, Herb. Kew. Nearest to the preceding species.

*Hirtella triandra*, Sw. Prodr. 57. Reis, 1,500 ft. (1448).

*Hirtella bracteata*, Mart. & Zucc. Abhand. Akad. Munch. x. 384? Guanai, 2,000 ft. (2492). Specimens insufficient for certain determination.

*Conepia glaucescens*, Spruce in Mart. Flor. Bras. xiv. (2) 49. Falls of Madeira, Brazil (2678).

*Prunus salicifolia*, HBK. Nov. Gen. iv. 241. Near La Paz, 10,000 ft. (480).

*Quillaja Saponaria*, Mol. Chil. 354. Sorata, 8,000 ft. (728).

*Rubus megallococcus*, Focke, Abhand. Nat. Gesell. Bremen, iv. 157. Unduavi, 8,000 ft. (470 and 472). The same as Mandon's 662.

*Rubus Boliviensis*, Focke, l. c. 158. Same locality (468). Distributed as *R. Sellowii*, C. & S.? The same as Mandon's 656.

*Rubus Bogotensis*, HBK. l. c. vi. 220. Unduavi, 8,000 ft. (474). Possibly this may be an extreme form of *R. floribundus*.

*Rubus roseus*, Poir. Dict. vi. 237. Unduavi, 8,000 ft. (473 and 469). The same as Mandon's Nos. 659 in part and 661.

*Rubus glaucus*, Benth. Pl. Hartw. 173. Yungas, 6,000 ft. (471).

*RUBUS RUSBYI*, spec. nova. Ascendens; tomentosus; folia integra, petiolata, serrata, utrinque tomentosa; stipulis latis,

- ovatis; ramulis, petiolisque minute recurvo-aculeatis; floribus solitariis, 2 cm. latis; sepalis ovatis, acutis, 8 cm. longis, dense tomentosus. Planta 50 cm. alta; folia 4 cm. longa, 2 cm. lata. Next to *R. Loxensis*, Benth. Unduavi, 10,000 ft. (2508).
- Fragaria Chilensis*, Ehrh. Beitr. vii. 26. Near La Paz, 10,000 ft. (479).
- Alchemilla aphanoides*, Mutis. in Linn. f. Suppl. 122. Unduavi, 8,000 ft. (851).
- Alchemilla hirsuta*, HBK. l. c. 226. Sorata, 13,000 ft. (477).
- Alchemilla tripartita*, R. & P. Fl. Per. i. 68. Unduavi, 10,000 ft. (476); 8,000 ft. (849 and 850, the latter a large, strong form).
- Alchemilla pectinata*, HBK. l. c. 226. Unduavi, 10,000 ft. (475).
- Alchemilla pinnata*, R. & P. l. c. 69. Near La Paz, 10,000 ft. (2551).
- Alchemilla*, probably a reduced form of *A. hirsuta*. Sorata, 13,000 ft. (1855). The same in Herb. Kew, coll. by Jameson in Columbia.
- Acæna ovalifolia*, R. & P. l. c. 67. Unduavi, 8,000 ft. (467).
- Acæna cylindrostachya*, R. & P. l. c. 68. Near La Paz, 10,000 ft. (466).
- Poterium Sanguisorba*, L. Sp. Pl. 1411. Near La Paz, 10,000 ft. (1926). Introduced from Europe.
- OSTEOMELES PERNETTYOIDES (Wedd.) (*Hesperomeles pernettyoides*, Wedd. Chlor. And. ii. 231). Sorata, 8,000 ft. (1787). Unduavi, 10,000 ft. (2039).

## SAXIFRAGÆ.

- Escallonia rubra*, Pers. Ench. i. 235. Valparaiso, Chili (1024).
- Escallonia revoluta*, (R. & P). Pers. l. c. Same locality (487 and 1772).
- PHYLLONOMA INTEGERRIMA (Turcz.) (*Dulongia integerrima*, Turcz. Bull. Soc. Moscou, 1858, 454). Mapiri, 5,000 ft. (2521). A form with serrate leaves. I am obliged to Mr. N. E. Brown for putting me on the track of this plant.
- Weinmannia hirtella*, HBK. l. c. 56. Mapiri, 10,000 ft. (2040).
- Weinmannia elliptica*, HBK. l. c. 50. Yungas, 6,000 ft. (535); Unduavi, 8,000 ft. (536). Some of these specimens were distributed as *W. subsessilifolia*, R. & P.
- Ribes albiflorum*, R. & P. Fl. Per. iii. 12. Unduavi, 8,000 ft. (478).

## HALORACEÆ.

*Gunnera scabra*, R. & P. l. c. t. 44. Unduavi, 8,000 ft. (624).

## COMBRETACEÆ.

*Combretum Jacquinii*, Griseb. Fl. Brit. West Indies, 275. Junction of the Beni and Madre de Dios (721).

*Combretum Læflingii*, Eichl. in Mart. Fl. Bras. xiv. (2) 110. Guanai, 2,000 ft. (622), and (2635?) the latter in fruit only.

*Combretum Aubletii*, DC. Prodr. iii. 19. Beni River (1773).

*Combretum*, sp. undetermined. Guanai, 2,000 ft. (623).

## MYRTACEÆ.

*Eucalyptus capitellatus*, Sm. Bot. Nov. Holl. 42. Near Valparaiso (600). Cultivated.

*Calyptranthes*, a species near *C. Spruceana*, Berg. Guanai, 2,000 ft. (1226). Not at Kew, and perhaps undescribed. Collected only in fruit.

*Psidium pomiferum*, L. Sp. Pl. 672. Beni River (2083).

*Psidium polycarpon*, Lamb. Trans. Linn. Soc. xi. 231. Yungas, 4,000 ft. (2460).

*Psidium*, sp. Falls of Madeira, Brazil, (2084).

*Psidium aromaticum*, Aubl. Guian. i. 485. Falls of Madeira (2085). Compared with Schomburgh's No. 355, Herb. Kew, and appears not distinct.

*Myrtus microphylla*, Humb. & Bonpl. Pl. Æquin. t. 4. Sorata, 10,000 ft. (2042).

*Myrcia lanceolata*, Camb. in St. Hil. Fl. Bras. Merid. ii. 236. Yungas, 6,000 ft. (605 and 1447).

*Myrcia Selloana*, Berg. in Mart. Flor. Bras. xiv. (1) 197. Falls of the Madeira (585); Junction of the Beni and Madre de Dios, (592) and 586).

*Myrcia velutina*, Berg. l. c. 182. Falls of the Madeira (590).

*Myrcia Paivæ*, Berg. l. c. 179. Yungas, 4,000 ft. (598). The same as Spruce No. 486.

*Myrcia anacardiæfolia*, Berg. Gardn. Lond. Journ. Bot. ii. 354. Falls of the Madeira (2683).

*Myrcia Berberis*, DC. Prodr. iii. 254? Falls of the Madeira (587).